Gene Cluster Activation in a Bacterial Symbiont Leads to Halogenated Angucyclic Maduralactomycins and Spirocyclic Actinospirols.

Guo H, Schwitalla JW, Benndorf R, Baunach M, Steinbeck C, Görls H, de Beer ZW, Regestein L, Beemelmanns C (2020) Gene Cluster Activation in a Bacterial Symbiont Leads to Halogenated Angucyclic Maduralactomycins and Spirocyclic Actinospirols. *Org Lett* 22(7), 2634-2638. <u>PubMed</u>

ILRS Authors

René Benndorf Jan Schwitalla

Projects

Investigation of secondary metabolites from insect-associated microbes and their contribution to insect homeostasis and defense Details

Metabolomic and transcriptomic analysis of the defensive role of Actinobacteria within the fungusgrowing termite system Details

Abstract

Growth from spores activated a biosynthetic gene cluster in *Actinomadura* sp. RB29, resulting in the identification of two novel groups of halogenated polyketide natural products, named maduralactomycins and actinospirols. The unique tetracyclic and spirocyclic structures were assigned based on a combination of NMR analysis, chemoinformatic calculations, X-ray crystallography, and 13C labeling studies. On the basis of HRMS2 data, genome mining, and gene expression studies, we propose an underlying noncanonical angucycline biosynthesis and extensive post-polyketide synthase (PKS) oxidative modifications.

Identifier

doi: 10.1021/acs.orglett.0c00601 PMID: 32193935