

# ***Nocardia macrotermitis* sp. nov. and *Nocardia aurantia* sp. nov., isolated from the gut of the fungus-growing termite *Macrotermes natalensis*.**

Benndorf R, Schwitalla JW, Martin K, de Beer ZW, Vollmers J, Kaster AK, Poulsen M, Beemelmanns C (2020) *Nocardia macrotermitis* sp. nov. and *Nocardia aurantia* sp. nov., isolated from the gut of the fungus-growing termite *Macrotermes natalensis*. *Int J Syst Evol Microbiol* , [PubMed](#)

## **ILRS Authors**

[René Benndorf](#) [Jan Schwitalla](#)

## **Projects**

Investigation of secondary metabolites from insect-associated microbes and their contribution to insect homeostasis and defense

[Details](#)

Metabolomic and transcriptomic analysis of the defensive role of Actinobacteria within the fungus-growing termite system

[Details](#)

## **Abstract**

The taxonomic positions of two novel aerobic, Gram-stain-positive Actinobacteria, designated RB20T and RB56T, were determined using a polyphasic approach. Both were isolated from the fungus-farming termite *Macrotermes natalensis*. Results of 16S rRNA gene sequence analysis revealed that both strains are members of the genus *Nocardia* with the closest phylogenetic neighbours *Nocardia miyunensis* JCM12860T (98.9 %) and *Nocardia nova* DSM44481T (98.5 %) for RB20T and *Nocardia takedensis* DSM 44801T (98.3 %), *Nocardia pseudobrasiliensis* DSM 44290T (98.3 %) and *Nocardia rayongensis* JCM 19832T (98.2 %) for RB56T. Digital DNA-DNA hybridization (DDH) between RB20T and *N. miyunensis* JCM12860T and *N. nova* DSM 44481T resulted in similarity values of 33.9 and 22.0 %, respectively. DDH between RB56T and *N. takedensis* DSM44801T and *N. pseudobrasiliensis* DSM44290T showed similarity values of 20.7 and 22.3 %, respectively. In addition, wet-lab DDH between RB56T and *N. rayongensis* JCM19832T resulted in 10.2 % (14.5 %) similarity. Both strains showed morphological and chemotaxonomic features typical for the genus *Nocardia*, such as the presence of meso-diaminopimelic acid (A2pm) within the cell wall, arabinose and galactose as major sugar components within whole cell-wall hydrolysates, the presence of mycolic acids and major phospholipids (diphosphatidylglycerol, phosphatidylethanolamine, phosphatidylinositol), and the predominant menaquinone MK-8 (H4, ω-cyclo). The main fatty acids for both strains were hexadecanoic acid (C16 : 0), 10-methyloctadecanoic acid (10-methyl C18 : 0) and cis-9-octadecenoic acid (C18 : 1 ω9c). We propose two novel species within the genus *Nocardia*: *Nocardia macrotermitis* sp. nov. with the type strain RB20T (=VKM Ac-2841T=NRRL B65541T) and *Nocardia aurantia* sp. nov. with the type strain RB56T (=VKM Ac-2842T=NRRL B65542T).

**Identifier**

doi: 10.1099/ijsem.0.004398 PMID: 32815801